

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human MX1 protein, His tag, Unconjugated GTX00244-PRO

Artikelname	Human MX1 protein, His tag, Unconjugated
Artikelnummer	GTX00244-PRO
Hersteller Artikelnummer	GTX00244-pro
Alternativnummer	GTX00244-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	4599
UniProt	P20591
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Ser80~Leu342 (NP_001138397.1)

Anwendungsbeschreibung

Interferon-induced GTP-binding protein Mx1 is a protein that in humans is encoded by the MX1 gene. In mouse, the interferon-inducible Mx protein is responsible for a specific antiviral state against influenza virus infection. The protein encoded by this gene is similar to the mouse protein as determined by its antigenic relatedness, induction conditions, physicochemical properties, and amino acid analysis. This cytoplasmic protein is a member of both the dynamin family and the family of large GTPases. Besides, Tubulin Beta (TUBb) has been identified as an interactor of MX1, thus a binding ELISA assay was conducted to detect the interaction of recombinant human MX1 and recombinant human TUBb. Briefly, MX1 was diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to TUBb-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-MX1 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of MX1 and TUBb was in a dose dependent manner.