

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## **Product Datasheet**

## Human Creatine kinase (brain) protein, His tag, Unconjugated GTX00246-PRO

Artikelname	Human Creatine kinase (brain) protein, His tag, Unconjugated
Artikelnummer	GTX00246-PRO
Hersteller Artikelnummer	GTX00246-pro
Alternativnummer	GTX00246-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	1152
UniProt	P12277
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Lys11~Leu367 (NP_001814.2)

Anwendungsbeschreibung

Creatine Kinase, Brain (CKB) also known as CK-BB is a creatine kinase. CKB, consists of a homodimer of two identical brain-type CK-B subunits, is a cytoplasmic enzyme involved in cellular energy homeostasis, with certain fractions of the enzyme being bound to cell membranes, ATPases, and a variety of ATP-requiring enzymes in the cell. Besides, Creatine Kinase, Muscle (CKM) has been identified as an interactor of CKB, thus a binding ELISA assay was conducted to detect the interaction of recombinant human CKB and recombinant human CKM. Briefly, CKB were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to CKM-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-CKB pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of CKB and CKM was in a dose dependent manner.