

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human CD14 protein, His tag (active), Unconjugated GTX00249-PRO

Artikelname	Human CD14 protein, His tag (active), Unconjugated
Artikelnummer	GTX00249-PRO
Hersteller Artikelnummer	GTX00249-pro
Alternativnummer	GTX00249-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	929
UniProt	P08571
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Asp125~Leu288 (NP_000582.1)

Anwendungsbeschreibung

Cluster Of Differentiation 14 (CD14), also known as CD14, is a component of the innate immune system. CD14 acts as a co-receptor (along with the Toll-like receptor TLR 4 and MD-2) for the detection of bacterial lipopolysaccharide (LPS). CD14 can bind LPS only in the presence of lipopolysaccharide-binding protein (LBP). Although LPS is considered its main ligand, CD14 also recognizes other pathogen-associated molecular patterns such as lipoteichoic acid. Besides, Lipopolysaccharide Binding Protein (LBP) has been identified as an interactor of CD14, thus a binding ELISA assay was conducted to detect the interaction of recombinant human CD14 and recombinant human LBP. Briefly, CD14 were diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to LBP-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-CD14 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of CD14 and LBP was in a dose dependent manner. CD14 also can enhance LPS-stimulated IL-8 secretion by THP-1 human acute monocytic leukemia cells. Therefore, THP-1 cells were cultured in 24 well plates at a concentration of 1×10^6 cells/ml and stimulated by LPS (1µg/ml), then added of different concentrations of recombinant human CD14 (1µg/ml, 5µg/ml). After 24h later, the production of IL8 was determined in the supernatants by cytokine specific ELISA. When the concentration of CD14 was 5µg/ml, the secretion LPS-stimulated IL-8 can be significantly increased.