

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Human FGF23 protein, His tag, Unconjugated GTX00257-PRO

Artikelname	Human FGF23 protein, His tag, Unconjugated
Artikelnummer	GTX00257-PRO
Hersteller Artikelnummer	GTX00257-pro
Alternativnummer	GTX00257-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	<a href="#">8074</a>
UniProt	<a href="#">Q9GZV9</a>
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Asp79~Arg160 (NP_065689.1)

Anwendungsbeschreibung

Fibroblast growth factor 23 or FGF23 is a member of the fibroblast growth factor (FGF) family which is responsible for phosphate and vitamin D metabolism. The main function of FGF23 seems to be regulation of phosphate concentration in plasma. FGF23 decreases the reabsorption and increases excretion of phosphate and suppress 1-alpha-hydroxylase, reducing its ability to activate vitamin D and subsequently impairing calcium absorption. Besides, Fibroblast Growth Factor Receptor 2 (FGFR2) has been identified as an interactor of FGF23, thus a binding ELISA assay was conducted to detect the interaction of recombinant human FGF23 and recombinant human FGFR2. Briefly, FGF23 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to FGFR2-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-FGF23 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of FGF23 and FGFR2 was in a dose dependent manner.