

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human SEMA3F protein, His tag (active), Unconjugated GTX00267-PRO

Artikelname	Human SEMA3F protein, His tag (active), Unconjugated
Artikelnummer	GTX00267-PRO
Hersteller Artikelnummer	GTX00267-pro
Alternativnummer	GTX00267-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	6405
UniProt	Q13275
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Val350~Leu483 (NP_001305729.1)

Anwendungsbeschreibung	<p>Semaphorin-3F (SEMA3F) is a member of semaphorins family. All the family members have a secretion signal, a 500-amino acid sema domain, and 16 conserved cysteine residues. Semaphorins not only guide axons in development, but also have major roles in immune function (classes 4, 6, and 7) and the development of bones. Class 3 semaphorins are one of the most versatile semaphorin classes, in which Sema3a is the most studied. Besides, SEMA3F is deleted or downregulated in many metastatic tumors. To test the effect of SEMA3F on inhibit cell proliferation, human lung carcinoma cells A549 were seeded into triplicate wells of 96-well plates at a density of 5000 cells/well and allowed to attach, replaced with serum-free overnight, then the medium was replaced with 2% serum standard DMEM prior to the addition of various concentrations of recombinant human SEMA3F. After incubated for 48h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 μl of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37C. Proliferation of A549 cells after incubation with SEMA3F for 48h observed by inverted microscope. Cell viability was assessed by CCK-8 assay after incubation with recombinant SEMA3F for 48h. And SEMA3F significantly increased cell viability of A549 cells.</p>
------------------------	---