

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse S100 beta protein, His tag, Unconjugated GTX00294-PRO

Artikelname	Mouse S100 beta protein, His tag, Unconjugated
Artikelnummer	GTX00294-PRO
Hersteller Artikelnummer	GTX00294-pro
Alternativnummer	GTX00294-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Mouse
Konjugation	Unconjugated
NCBI	20203
UniProt	P50114
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	Full length protein, N-terminal His-Tag, Met1~Glu92 (NP_033141.1)

Anwendungsbeschreibung

Protein S100B is a member of the S100 family. S100 proteins are EFhand calcium-binding proteins and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. Experimental results suggest that the receptor for advanced glycation end products (RAGE) plays important roles in mediating S100 protein-induced cellular signaling. Besides, mouse RAGE shares similarities with human RAGE in amino acids sequence with the identity of 78.2% . Thus a binding ELISA assay was conducted to detect the interaction of recombinant mouse S100B and recombinant human RAGE. Briefly, S100B were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to RAGE-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-S100B mAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of of S100B and RAGE was in a dose dependent manner.