

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse IL2 Receptor alpha protein, His tag, Unconjugated GTX00298-PRO

Artikelname	Mouse IL2 Receptor alpha protein, His tag, Unconjugated
Artikelnummer	GTX00298-PRO
Hersteller Artikelnummer	GTX00298-pro
Alternativnummer	GTX00298-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Mouse
Konjugation	Unconjugated
NCBI	16184
UniProt	P01590
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Glu22~Lys236 (NP_032393.3)

Anwendungsbeschreibung

IL2RA (Interleukin-2 receptor alpha chain), also known as Tac antigen and as CD25, was initially identified as a 55kDa membrane glycoprotein which is capable of binding IL 2. Besides, mouse IL2 shares 75.1% AA sequence identity with rat IL2, suggesting the exist of cross-species activity. Thus we have conducted a binding ELISA assay to detect the interaction of recombinant rat IL2RA with recombinant mouse IL2. Briefly, IL2RA were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 μl were then transferred to IL2-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-IL2RA pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 μ l stop solution to the wells and read at 450nm immediately. The binding activity of IL2RA with IL2 was in a dose dependent manner.