

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## **Product Datasheet**

## Mouse iNOS protein, His tag, Unconjugated GTX00325-PRO

Artikelname	Mouse iNOS protein, His tag, Unconjugated
Artikelnummer	GTX00325-PRO
Hersteller Artikelnummer	GTX00325-pro
Alternativnummer	GTX00325-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Mouse
Konjugation	Unconjugated
NCBI	18126
UniProt	P29477
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Asp43~Thr213 (NP_001300850.1)

Anwendungsbeschreibung

Nitric oxide synthase 2, inducible (NOS2) is a member of Nitric oxide synthases (NOSs) family. Nitric oxide synthases (NOSs) are a family of enzymes catalyzing the production of nitric oxide (NO) from Larginine. NO is an important cellular signaling molecule. It helps modulate vascular tone, insulin secretion, airway tone, and peristalsis, and is involved in angiogenesis and neural development. Besides, Ubiquitin Carboxyl Terminal Hydrolase L5 (UCHL5) has been identified as an interactor of NOS2, thus a binding ELISA assay was conducted to detect the interaction of recombinant mouse NOS2 and recombinant mouse UCHL5. Briefly, NOS2 was diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100  $\mu$ l were then transferred to UCHL5-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-NOS2 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 μl stop solution to the wells and read at 450nm immediately. The binding activity of NOS2 and UCHL5 was in a dose dependent manner.