

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit CXCL8 / IL8 protein, His tag (active), Unconjugated GTX00339-PRO

Artikelname	Rabbit CXCL8 / IL8 protein, His tag (active), Unconjugated
Artikelnummer	GTX00339-PRO
Hersteller Artikelnummer	GTX00339-pro
Alternativnummer	GTX00339-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Rabbit
Konjugation	Unconjugated
NCBI	100009129
UniProt	P19874
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Ala23~Ser101 (NP_001075762.1)

undertaken to detect the chemotactic effect of IL8 on the human T-lymphocyte leukemia cell line Jurkat. Briefly, Jurkat cells were seeded into the upper chambers (100 μ l cell suspension, 10 6 cells/ml in RPMI-1640 with FBS free) and IL8 (0. 1 ng/ml, 1 ng/ml and 10 ng/ml diluted separately in serum free RPMI-1640) was added in lower chamber with a polycarbonate filter (8um pore size) used to separate the two compartments. After incubation at 37C with 5% CO2 for 1h, the filter was removed, then cells in low chamber were

observed by inverted microscope at low magnification (*100) and the number of migrated cells were counted at high magnification (*400) randomly (five fields for each filter). The migrated Jurkat cells

in low chamber at low magnification (*100) randomly, and the

IL8 occurs at $1\sim10$ ng/ml.

migrated cells were counted at high magnification (*400). IL8 is able to induce migration of Jurkat cells, and the optimum chemotaxis of

Interleukin 8 (IL8 or chemokine (C-X-C motif) ligand 8, CXCL8) is a chemokine produced by macrophages and other cell types such as epithelial cells, airway smooth muscle cells and endothelial cells. Thus, chemotaxis assay used 24-well microchemotaxis system was

Anwendungsbeschreibung