

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human MCP1 / CCL2 protein, His tag (active), Unconjugated GTX00415-PRO

| Artikelname | Human MCP1 / CCL2 protein, His tag (active), Unconjugated |
|--------------------------|---|
| Artikelnummer | GTX00415-PRO |
| Hersteller Artikelnummer | GTX00415-pro |
| Alternativnummer | GTX00415-PRO-10 |
| Hersteller | GeneTex |
| Kategorie | Proteine/Peptide |
| Applikation | FA |
| Spezies Reaktivität | Human |
| Konjugation | Unconjugated |
| NCBI | 6347 |
| UniProt | P13500 |
| Puffer | Reconstitute with 20mM Tris (pH8.0) and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris (pH8.0), 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300. |
| Expression System | E. coli |
| Formulierung | Lyophilized powder |
| Sequenz | N-terminal His-Tag, Gln24~Thr99 (NP_002973.1) |

Anwendungsbeschreibung

MCP-1 (monocyte chemoattractant protein 1), also known as CCL2 (C-C motif chemokine 2), is a small cytokine that belongs to the CC chemokine family. MCP-1 has been described as a chemoattractant for monocytes and proven to be able to induce chemotactic migration of THP-1 cells. Therefore, chemotaxis assay used 24-well microchemotaxis system was undertaken to detect the chemotactic effect of MCP-1 on the human monocytic cell line THP-1. Briefly, THP-1 cells were seeded into the upper chambers (100 µl cell suspension, 106 cells/ml in RPMI-1640 with 0.5% FBS) and MCP-1 (25 and 50 ng/ml diluted separately in serum free RPMI-1640) was added in lower chamber with a polycarbonate filter (8 µm pore size) used to separate the two compartments. After incubation at 37C with 5% CO2 for 3h, the filter was removed, then cells in low chamber were observed by inverted microscope at low magnification (*100) and the number of migrated cells were counted at high magnification (*400) randomly (five fields for each filter). MCP-1 is able to induce migration of THP-1 cells, and the optimum chemotactic concentration of MCP-1 is about 25 ng/ml.