

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human EPO protein, His and GST tag (active), Unconjugated GTX00422-PRO

Artikelname	Human EPO protein, His and GST tag (active), Unconjugated
Artikelnummer	GTX00422-PRO
Hersteller Artikelnummer	GTX00422-pro
Alternativnummer	GTX00422-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	ELISA, FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	2056
UniProt	P01588
Puffer	Reconstitute with 20mM Tris (pH8.0) and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris (pH8.0), 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag and GST-Tag, Ala28~Arg193 (NP_000790.2)

Anwendungsbeschreibung

Erythropoietin (EPO), also known as hematopoietin or hemopoietin, is a glycoprotein cytokine secreted by the kidney in response to cellular hypoxia, it stimulates red blood cell production (erythropoiesis) in the bone marrow. Erythropoietin is an essential hormone for red blood cell production. EPO can cooperate with various other growth factors involved in the development of erythroid lineage from multipotent progenitors. To test the effect of EPO on cell proliferation, TF-1 cells were seeded into triplicate wells of 96-well plates at a density of 5000 cells/well with 1% serum standard 1640 including various concentrations of recombinant human EPO. After incubated for 72h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 µl of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37C. Proliferation of TF-1 cells after incubation with EPO for 72h observed by inverted microscope, and cell viability was assessed by CCK-8 (Cell Counting Kit-8) assay after incubation with recombinant EPO for 72h. EPO significantly increased cell viability of TF-1 cells.