

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Beta Catenin (p120)(Rabbit PAb), CF740 conjugate, 0.1mg/mL, Clone: [PAb], CF 740, Monoclonal BOT-BNC740376-500**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Beta Catenin (p120)(Rabbit PAb), CF740 conjugate, 0.1mg/mL, Clone: [PAb], CF 740, Monoclonal                                                                                                                                                                  |
| Artikelnummer            | BOT-BNC740376-500                                                                                                                                                                                                                                             |
| Hersteller Artikelnummer | BNC740376-500                                                                                                                                                                                                                                                 |
| Alternativnummer         | BOT-BNC740376-500-500UL                                                                                                                                                                                                                                       |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Rabbit                                                                                                                                                                                                                                                        |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | FC, IHC, WB                                                                                                                                                                                                                                                   |
| Spezies Reaktivität      | Human, Mouse                                                                                                                                                                                                                                                  |
| Immunogen                | A synthetic peptide from the middle of beta-Catenin (p120) protein                                                                                                                                                                                            |
| Konjugation              | CF 740                                                                                                                                                                                                                                                        |
| Produktbeschreibung      | Beta-catenin associates with the cytoplasmic portion of E-cadherin. The catenin/cadherin complexes play an important role mediating cellular adhesion, including adherens junctions. Beta-catenin is involved in the Wnt signaling pathway as well as othe... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [PAb]                                                                                                                                                                                                                                                         |

|                        |                                                                                                                                                                                                                                                                                                                                                                           |
|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | 92 kDa                                                                                                                                                                                                                                                                                                                                                                    |
| UniProt                | <a href="#">P35222</a>                                                                                                                                                                                                                                                                                                                                                    |
| Puffer                 | PBS, 0.1% rBSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                               |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                    |
| Anwendungsbeschreibung | Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |