

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker)(MOC-31), CF740 conjugate, 0.1mg/mL, Clone: [MOC-31], CF 740, Mouse, Monoclonal BOT-BNC740380-500

Artikelname	Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker)(MOC-31), CF740 conjugate, 0.1mg/mL, Clone: [MOC-31], CF 740, Mouse, Monoclonal
Artikelnummer	BOT-BNC740380-500
Hersteller Artikelnummer	BNC740380-500
Alternativnummer	BOT-BNC740380-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Neuraminidase treated GLS-1 human small cell lung carcinoma cells
Konjugation	CF 740
Produktbeschreibung	The epitope of this antibody is located in the first EGF-like repeat domain (EGF1) between amino acids 27-59 of Ep-CAM. EGP40 is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cel...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[MOC-31]
Molekulargewicht	40-43 kDa
UniProt	P16422
Puffer	PBS, 0.1% rBSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT Western blot: 1-2 ug/mL Does not react with rat Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user