

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

NKX2.2 (Neuroendocrine & Ewing s Sarcoma Marker) (NX2/1523), CF740 conjugate, 0.1mg/mL, Clone: [NX2/1523], CF 740, Mouse, Monoclonal BOT-BNC741523-100

Artikelname	NKX2.2 (Neuroendocrine & Ewing s Sarcoma Marker) (NX2/1523), CF740 conjugate, 0.1mg/mL, Clone: [NX2/1523], CF 740, Mouse, Monoclonal
Artikelnummer	BOT-BNC741523-100
Hersteller Artikelnummer	BNC741523-100
Alternativnummer	BOT-BNC741523-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Human recombinant NKX2.2 protein fragment (aa1-119) (exact sequence is proprietary)
Konjugation	CF 740
Produktbeschreibung	Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein s...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[NX2/1523]
Molekulargewicht	40-50 kDa
UniProt	O95096
Puffer	PBS, 0.1% rBSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user