

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Tyrosinase (Melanoma Marker) (TYR/2024R), CF740 conjugate, 0.1mg/mL, Clone: [TYR/2024R], CF 740, Rabbit, Monoclonal BOT-BNC742024-500

Artikelname	Tyrosinase (Melanoma Marker) (TYR/2024R), CF740 conjugate, 0.1mg/mL, Clone: [TYR/2024R], CF 740, Rabbit, Monoclonal
Artikelnummer	BOT-BNC742024-500
Hersteller Artikelnummer	BNC742024-500
Alternativnummer	BOT-BNC742024-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human tyrosinase (TYR) protein
Konjugation	CF 740
Produktbeschreibung	This antibody recognizes a cluster of proteins between 70-80 kDa, identified as tyrosinase. Occasionally a minor band at 55 kDa is also detected. This MAb shows no cross-reaction with MAGE-1 and tyrosinase-related protein 1, TRP-1/gp75. Tyrosinase is...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[TYR/2024R]

Molekulargewicht	70-80 kDa
UniProt	P14679
Puffer	PBS, 0.1% rBSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 or 10 mM citrate buffer pH6.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user