

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

**DOG-1 / TMEM16A (Gastrointestinal Stromal Tumor Marker) (DG1/2564R),
CF740 conjugate, 0.1mg/mL, Clone: [DG1/2564R], CF 740, Rabbit,
Monoclonal
BOT-BNC742564-500**

Artikelname	DOG-1 / TMEM16A (Gastrointestinal Stromal Tumor Marker) (DG1/2564R), CF740 conjugate, 0.1mg/mL, Clone: [DG1/2564R], CF 740, Rabbit, Monoclonal
Artikelnummer	BOT-BNC742564-500
Hersteller Artikelnummer	BNC742564-500
Alternativnummer	BOT-BNC742564-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human DOG-1 protein fragment (aa 2-101) (exact sequence is proprietary)
Konjugation	CF 740
Produktbeschreibung	Expression of DOG-1 protein is elevated in the gastrointestinal stromal tumors (GIST s), c-kit signaling-driven mesenchymal tumors of the GI tract. DOG-1 is rarely expressed in other soft tissue tumors, which, due to appearance, may be difficult to d...
Klonalität	Monoclonal

Konzentration	0.1 mg/mL
Klon-Bezeichnung	[DG1/2564R]
Molekulargewicht	~114 kDa
UniProt	Q5XXA6
Puffer	PBS, 0.1% rBSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user