

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Myogenin (Skeletal Muscle Marker) (MYOG/2660), CF740 conjugate, 0.1mg/mL, Clone: [MYOG/2660], CF 740, Mouse, Monoclonal BOT-BNC742660-500

Artikelname	Myogenin (Skeletal Muscle Marker) (MYOG/2660), CF740 conjugate, 0.1mg/mL, Clone: [MYOG/2660], CF 740, Mouse, Monoclonal
Artikelnummer	BOT-BNC742660-500
Hersteller Artikelnummer	BNC742660-500
Alternativnummer	BOT-BNC742660-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human myogenin (MYOG) protein
Konjugation	CF 740
Produktbeschreibung	Myogenin is a member of the MyoD family of myogenic basic helix-loop-helix (bHLH) transcription factors that also includes MyoD, Myf-5, and MRF4 (also known as herculinor Myf-6). MyoD family members are expressed exclusively in skeletal muscle and pl...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[MYOG/2660]

Molekulargewicht	34 kDa
UniProt	P15173
Puffer	PBS, 0.1% rBSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user