

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Human IL-7 Protein, HEK293 Lysate RAY-230-10274-100

Artikelname	Recombinant Human IL-7 Protein, HEK293 Lysate
Artikelnummer	RAY-230-10274-100
Hersteller Artikelnummer	230-10274-100
Alternativnummer	RAY-230-10274-100
Hersteller	RayBiotech
Kategorie	Proteine/Peptide
Spezies Reaktivität	Human
Produktbeschreibung	Recombinant human Interleukin-7 overexpression cell lysate, derived from the transfected HEK293 cells (100 µg). Purchase will also include one vial of normal control HEK293 cell lysate (catalog . 230-10006) transfected with empty expression vector....
Konzentration	Determined by Pierce BCA protein assay kit
Molekulargewicht	Recombinant protein has a calculated molecular mass of 17. The actual molecular weight may increase differently due to the potential post-modifications (PTMs).
Tag	His
Expression System	HEK293 cells

Reinheit	<p>Unpurified cell lysate. HEK293 cells transfected with expression vectors harboring target gene were harvested and washed with PBS twice. The cell pastes were re-suspended with ice-cold PBS containing mammalian cell protease inhibitor cocktail and further lysed with freeze-thaw cycles. After clarifying with 20,000 g centrifugation at 4C for 30 min, the lysate was aliquoted, lyophilized, and stored at -80C immediately. Protein concentration was determined by BCA kit (Thermo Scientific, Inc.) using BSA as protein standard. The gene overexpression in lysate was confirmed by Western blotting using anti-His tag antibody and/or target-specific antibodies and the lysate derived from HEK293 cells transfected with the empty expression vector was used as a negative control.</p>
Formulierung	Lyophilized powder
Sequenz	Asp26-His177
Formel	Lyophilized from a 0.2 µm filtered solution in PBS containing mammalian cell protease inhibitor cocktail
Anwendungsbeschreibung	Briefly spin the vial and bring the contents to the bottom prior to opening. It is recommended to reconstitute at 0.5 - 1 mg/mL with sterile deionized water.