

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### TNFSF11 Antibody (Biotin), Goat, Polyclonal BYT-ORB1272305

|                          |   |
|--------------------------|---|
| Artikelname              | TNFSF11 Antibody (Biotin), Goat, Polyclonal   |
| Artikelnummer            | BYT-ORB1272305  |
| Hersteller Artikelnummer | orb1272305  |
| Alternativnummer         | BYT-ORB1272305-0.05   |
| Hersteller               | Biorbyt   |
| Wirt                     | Goat  |
| Kategorie                | Antikörper  |
| Applikation              | ELISA, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hsRANKL (Human sRANK-Ligand). |
| Konjugation              | Biotin  |
| Produktbeschreibung      | TNFSF11 Antibody (Biotin)...  |
| Klonalität               | Polyclonal  |
| Konzentration            | batch dependent   |
| NCBI                     | <a href="#">14788</a>   |
| UniProt                  | <a href="#">O14788</a>  |
| Formulierung             | Lyophilized   |

|                        |   |
|------------------------|---|
| Target-Kategorie       | TNFSF11   |
| Anwendungsbeschreibung | <p>Application Notes: ELISA:Sandwich:To detect hsRANKL by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.25 - 1.0 µg/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our Polyclonal Anti-Human sRANKL (XP-5274) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hsRANKL. Western Blot:To detect hsRANKL by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hsRANKL is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions</p> |