

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Human Tumor-associated calcium signal transducer 2 (TACSTD2), partial (Active) BYT-ORB1674363

Artikelname	Recombinant Human Tumor-associated calcium signal transducer 2 (TACSTD2), partial (Active)
Artikelnummer	BYT-ORB1674363
Hersteller Artikelnummer	orb1674363
Alternativnummer	BYT-ORB1674363-1,BYT-ORB1674363-100
Hersteller	Biorbyt
Kategorie	Proteine/Peptide
Produktbeschreibung	This Recombinant Human Tumor-associated calcium signal transducer 2 (TACSTD2), partial (Active) spans the amino acid sequence from region 31-274aa. Purity: Greater than 95% as determined by SDS-PAGE....
Molekulargewicht	30.3 kDa
UniProt	P09758
Puffer	Lyophilized from a 0.2 µm filtered 20 mM Tris-HCl, 0.5 M NaCl, 6% Trehalose, pH 8.0
Quelle	Homo sapiens (Human)
Reinheit	Greater than 95% as determined by SDS-PAGE.
Formulierung	Lyophilized powder

Sequenz	QDNCTCPTNKMTVCSPDGGPGGRCQCRA LGSGMAVDCSTLTSKCLLLKARMS APKNARTLVRPSEHALVDNDGLYDPDCDPEGRFKARQCNQTSVCWCVNSVG VRRTDKGDLSLRCDELVRTHHILIDLRHRPTAGAFNHSDLDAELRRLFRERYR LHPKFVA AVHYEQPTIQIELRQNTSQKAAGDVDIGDAAYYFERDIKGESLFQGR GGLDLRVRGEPLQVERTLIYYLDEIPPKFSMKRLT
Anwendungsbeschreibung	Biological Origin: Homo sapiens (Human). Biological Activity: Measured by its binding ability in a functional ELISA. Immobilized Human TROP2 at 2 µg/mL can bind Anti-TROP2 recombinant antibody, the EC50 is 0.9108-1.640 ng/mL. Application Notes: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference