

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Recombinant Human Proprotein convertase subtilisin/kexin type 9 (PCSK9) (D374Y), Biotinylated (Active) BYT-ORB1881659

Artikelname	Recombinant Human Proprotein convertase subtilisin/kexin type 9 (PCSK9) (D374Y), Biotinylated (Active)
Artikelnummer	BYT-ORB1881659
Hersteller Artikelnummer	orb1881659
Alternativnummer	BYT-ORB1881659-1,BYT-ORB1881659-100,BYT-ORB1881659-20
Hersteller	Biorbyt
Kategorie	Proteine/Peptide
Produktbeschreibung	This Recombinant Human Proprotein convertase subtilisin/kexin type 9 (PCSK9) (D374Y), partial, Biotinylated spans the amino acid sequence from region 31-692aa(D374Y). Purity: Greater than 95% as determined by SDS-PAGE....
Molekulargewicht	76.0 kDa
UniProt	<a href="#">Q8NBP7</a>
Puffer	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose, pH 8.0.
Quelle	Homo sapiens (Human)
Reinheit	Greater than 95% as determined by SDS-PAGE.
Formulierung	Lyophilized powder

Sequenz	QEDEDGDYEELVLALRSEEDGLAEAPEHGTTATFHRCADPWRLPGTYVVVL KEETHLSQSERTARRLQAQAARRGYLTKILHVFHGLLPGLVKMSGDLLELAL KLPHVDYIEEDSSVFAQSIPWNLERITPPRYRADEYQPPDGGSLVEVYLLDTSI QSDHREIEGRVMVTD FENVPEEDGTRFHRQASKCDSHGTHLAGVVSGRDAG VAKGASMRSLRVLNCOGKGT VSGTLIGLEFIRKSQLVQPVGPLVV
Anwendungsbeschreibung	<p>Biological Origin: Homo sapiens (Human). Biological Activity: Measured by its binding ability in a functional ELISA. Immobilized Anti-PCSK9 recombinant antibody at 2 µg/mL can bind Human PCSK9 protein. The EC50 is 7.450-8.692 ng/mL. ②Measured by its binding ability in a functional ELISA. Immobilized Human PCSK9 at 2 µg/mL on streptavidin coated plates can bind Anti-PCSK9 recombinant antibody. The EC50 is 0.3792-1.157 ng/mL. Application Notes: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference</p>