

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **TIGIT / VSTM3 / VSIG9 (Immune Checkpoint for Cancer) Antibody, IgG2b, Clone: [TIGIT/3106], Biotin, Mouse, Monoclonal BYT-ORB751152**

|                          |  |
|--------------------------|--|
| Artikelname              | TIGIT / VSTM3 / VSIG9 (Immune Checkpoint for Cancer) Antibody, IgG2b, Clone: [TIGIT/3106], Biotin, Mouse, Monoclonal |
| Artikelnummer            | BYT-ORB751152  |
| Hersteller Artikelnummer | orb751152  |
| Alternativnummer         | BYT-ORB751152-100  |
| Hersteller               | Biorbyt  |
| Wirt                     | Mouse  |
| Kategorie                | Antikörper   |
| Applikation              | ELISA, IHC   |
| Spezies Reaktivität      | Human  |
| Immunogen                | Recombinant fragment (around aa 22-141) of human TIGIT protein (exact sequence is proprietary)                       |
| Konjugation              | Biotin   |
| Produktbeschreibung      | Mouse monoclonal antibody against TIGIT conjugated to Biotin...  |
| Klonalität               | Monoclonal   |
| Konzentration            | Purified Ab conjugated to Biotin   |
| Klon-Bezeichnung         | [TIGIT/3106]   |
| Molekulargewicht         | 26kDa  |

|                        |   |
|------------------------|---|
| Isotyp                 | IgG2b   |
| UniProt                | <a href="#">Q495A1</a>  |
| Puffer                 | Antibody purified from Bioreactor Concentrate by Protein A/G and conjugated to Biotin. Prepared in 10mM PBS with 0.05% rAlbumin and 0.05% azide.  |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% rAlbumin & 0.05% azide. Also available WITHOUT rAlbumin & azide at 1.0mg/ml.  |
| Target-Kategorie       | TIGIT   |
| Application Verdünnung | Immunohistochemistry (Formalin-fixed) (2-4ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 C followed by cooling at RT for 20 minutes),Optimal dilution fo |
| Anwendungsbeschreibung | Application Notes: Positive Control: Human tonsil or prostate carcinoma   |