

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-MyD88 Monoclonal Antibody [2E9C2], IgG2b, Clone: [Clone: Clone: 2E9C2], Mouse BOB-M00025-2

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Anti-MyD88 Monoclonal Antibody [2E9C2], IgG2b, Clone: [Clone: Clone: 2E9C2], Mouse |
| Artikelnummer | BOB-M00025-2 |
| Hersteller Artikelnummer | M00025-2 |
| Alternativnummer | BOB-M00025-2-0.1MG |
| Hersteller | Boster Bio |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, IF, IHC-P, WB |
| Spezies Reaktivität | Human, Mouse |
| Immunogen | Mouse monoclonal MyD88 antibody was raised against a recombinant protein corresponding to amino acids 176 to 280 of human MyD88. |
| Produktbeschreibung | Boster Bio Anti-MyD88 Monoclonal Antibody [2E9C2] (Catalog M00025-2). Tested in ELISA, WB, IHC-P, IF applications. This antibody reacts with Human, Mouse.... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [Clone: Clone: 2E9C2] |

| | |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht | 33233 MW |
| Isotyp | IgG2b |
| UniProt | Q99836 |
| Puffer | MyD88 Monoclonal Antibody is supplied in PBS containing 0.02% sodium azide. |
| Formulierung | Liquid |
| Application Verdünnung | MyD88 antibody can be used for detection of MyD88 by Western blot at 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in mouse samples and Immunohi |