

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Human IGF1 protein, His tag (active), Unconjugated GTX00085-PRO**

Artikelname	Human IGF1 protein, His tag (active), Unconjugated
Artikelnummer	GTX00085-PRO
Hersteller Artikelnummer	GTX00085-pro
Alternativnummer	GTX00085-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	<a href="#">3479</a>
UniProt	<a href="#">P05019</a>
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	HEK293 cells
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, EFKLAAALEGGRKLEF+Gly49~Ala118 (NP_000609.1)

Anwendungsbeschreibung

insulin-like growth factor I (IGF1), is a hormone similar in molecular structure to insulin but have a much higher growth-promoting activity, it belongs to a family of proteins involved in mediating growth and development. It is reported that IGF1 induces the proliferation, migration, differentiation of a large types of cells including the MCF-7 breast cancer cell line. To test the effect of growth factors on proliferation, MCF-7 cells were seeded into triplicate wells of 96-well plates at a density of 2000 cells/well and allowed to attach overnight, then the medium was replaced with serum-free standard DMEM prior to the addition of various concentrations of IGF-1. After incubated for 72h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10  $\mu$ l of CCK-8 solution was added to each well of the plate, then measure the absorbance at 450nm using a microplate reader after incubating the plate for 1-4 hours at 37C. Cell proliferation of MCF-7 cells after incubation with IGF1 for 72h observed by inverted microscope.