

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human IL17 Receptor D protein, His tag, Unconjugated GTX00142-PRO

| | |
|--------------------------|---|
| Artikelname | Human IL17 Receptor D protein, His tag, Unconjugated |
| Artikelnummer | GTX00142-PRO |
| Hersteller Artikelnummer | GTX00142-pro |
| Alternativnummer | GTX00142-PRO-10 |
| Hersteller | GeneTex |
| Kategorie | Proteine/Peptide |
| Applikation | FA |
| Spezies Reaktivität | Human |
| Konjugation | Unconjugated |
| NCBI | 54756 |
| UniProt | Q8NFM7 |
| Puffer | Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300. |
| Expression System | E. coli |
| Formulierung | Lyophilized powder |
| Sequenz | N-terminal His-Tag, Thr157~Arg299 (NP_001305793.1) |

Anwendungsbeschreibung

IL17RD (interleukin-17 receptor D) is a membrane protein belonging to the IL17R protein family, acting as a component of the interleukin-17 receptor signaling complex. Knowing that the interaction between this protein and IL-17R does not require the interleukin, we have conducted a binding a binding ELISA assay to detect the interaction of recombinant human IL17RD with both recombinant human IL17RA and IL17. Briefly, IL17RD were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to IL17RA-coated or IL17-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-IL17RA pAb and anti-IL17 pAb separately, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution , wells were incubated 15-25 minutes at 37C. Finally, add 50 μ l stop solution to the wells and read at 450nm immediately. The binding activity of IL17RD with IL17RA and IL17 was shown in images respectively and this effect was in a dose dependent manner.