

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Human JAK2 protein, His tag, Unconjugated GTX00152-PRO

Artikelname	Human JAK2 protein, His tag, Unconjugated
Artikelnummer	GTX00152-PRO
Hersteller Artikelnummer	GTX00152-pro
Alternativnummer	GTX00152-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	<a href="#">3717</a>
UniProt	<a href="#">O60674</a>
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Asn508~Ala800 (NP_001309123.1)

#### Anwendungsbeschreibung

JAK2 (Tyrosine-protein kinase JAK2) is a tyrosine kinase involved in various processes such as cell growth, development, differentiation or histone modifications. JAK2 is considered to associate with some type I receptors, including EPOR (Erythropoietin receptor), therefore participates in cellular signal transduction. Thus a binding ELISA assay was conducted to detect the interaction of recombinant human JAK2 and recombinant human EPOR. Briefly, JAK2 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100  $\mu$ l JAK2 were then transferred to EPOR-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-JAK2 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50  $\mu$ l stop solution to the wells and read at 450nm immediately. The binding activity of JAK2 and EPOR was in a dose dependent manner.