

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human ALDH1A1 protein, His tag, Unconjugated GTX00155-PRO

Artikelname	Human ALDH1A1 protein, His tag, Unconjugated
Artikelnummer	GTX00155-PRO
Hersteller Artikelnummer	GTX00155-pro
Alternativnummer	GTX00155-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	216
UniProt	P00352
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Ser2~Ser501 (NP_000680.2)

Anwendungsbeschreibung

ALDH1A1 (Retinal dehydrogenase 1) which belongs to the aldehyde dehydrogenase family, is the next enzyme after alcohol dehydrogenase in the major pathway of alcohol metabolism. ALDH1A1 converts retinaldehyde to retinoic acid. IL21 (interleukin 21) was identified as an interactor of ALDH1A1 through Affinity Capture-MS. Thus a binding ELISA assay was conducted to detect the interaction of recombinant human ALDH1A1 and recombinant human IL21. Briefly, ALDH1A1 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl ALDH1A1 were then transferred to IL21-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-ALDH1A1 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of ALDH1A1 and IL21 was in a dose dependent manner.