

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human CD73 protein, His tag, Unconjugated GTX00159-PRO

Artikelname	Human CD73 protein, His tag, Unconjugated
Artikelnummer	GTX00159-PRO
Hersteller Artikelnummer	GTX00159-pro
Alternativnummer	GTX00159-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	4907
UniProt	P21589
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Leu29~Thr500 (NP_001191742.1)

Anwendungsbeschreibung

5-Nucleotidase, Ecto (NT5E), also known as ecto-5-nucleotidase or CD73, is an enzyme catalyzing the hydrolysis of nucleoside-5-monophosphates to nucleosides and inorganic phosphate. The enzyme is a dimer composed of 2 identical 70kD subunits bound by a glycosyl phosphatidyl inositol linkage to the external face of the plasma membrane. NT5E is a marker of lymphocyte differentiation that has functions independent of its catalytic activity, such as T-cell activation and cell-cell adhesion. Other forms of 5-prime nucleotidase exist in the cytoplasm and lysosomes and can be distinguished from NT5E by their substrate affinities, requirement for divalent magnesium ion, activation by ATP, and inhibition by inorganic phosphate. The enzyme is widely distributed in human and animal tissues. Besides, AF4/FMR2 Family, Member 1 (AFF1) has been identified as an interactor of NT5E thus a binding ELISA assay was conducted to detect the interaction of recombinant human NT5E and recombinant human AFF1. Briefly, NT5E were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to AFF1-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-NT5E pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 μ l stop solution to the wells and read at 450nm immediately. The binding activity of NT5E and AFF1 was in a dose dependent manner.