

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human Arylsulfatase F protein, His tag, Unconjugated GTX00162-PRO

Artikelname	Human Arylsulfatase F protein, His tag, Unconjugated
Artikelnummer	GTX00162-PRO
Hersteller Artikelnummer	GTX00162-pro
Alternativnummer	GTX00162-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	416
UniProt	P54793
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	HEK293 cells
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, His23~Asp300 (NP_001188467.1)

Anwendungsbeschreibung

ARSF (Arylsulfatase F) is a member of the sulfatase gene family, and more specifically, the arylsulfatase subfamily. Sulfatases are essential for the correct composition of bone and cartilage matrix. The activity of this protein, unlike that of arylsulfatase E, is not inhibited by warfarin. Besides, EP300 (E1A binding protein p300) has been identified as an interactor of ARSF, thus a binding ELISA assay was conducted to detect the interaction of recombinant human ARSF and recombinant human EP300. Briefly, ARSF were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to EP300-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-ARSF pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of ARSF and EP300 was in a dose dependent manner.