

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Human beta Tubulin I protein, His tag, Unconjugated GTX00175-PRO

Artikelname	Human beta Tubulin I protein, His tag, Unconjugated
Artikelnummer	GTX00175-PRO
Hersteller Artikelnummer	GTX00175-pro
Alternativnummer	GTX00175-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	<a href="#">203068</a>
UniProt	<a href="#">P07437</a>
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Val170~Val419 (NP_001280141.1)

#### Anwendungsbeschreibung

Tubulin Beta (TUBb) in molecular biology can refer either to the tubulin protein superfamily of globular proteins, or one of the member proteins of that superfamily. It participate in many essential cellular processes, including mitosis. alpha- and beta-tubulins polymerize into microtubules, a major component of the eukaryotic cytoskeleton. Both alpha and beta tubulins have a mass of around 50kDa and are thus in a similar range compared to actin with ~42kDa. TUBb is one of six members of the tubulin superfamily, which binding drugs to kill cancerous cells by inhibiting microtubule dynamics. Besides, Myxovirus Resistance 1 (MX1) has been identified as an interactor of TUBb, thus a binding ELISA assay was conducted to detect the interaction of recombinant human TUBb and recombinant human MX1. Briefly, TUBb were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to MX1-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-TUBb pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of TUBb and MX1 was in a dose dependent manner.