

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Human CD66e protein, His tag, Unconjugated GTX00181-PRO

|                          |   |
|--------------------------|---|
| Artikelname              | Human CD66e protein, His tag, Unconjugated  |
| Artikelnummer            | GTX00181-PRO  |
| Hersteller Artikelnummer | GTX00181-pro  |
| Alternativnummer         | GTX00181-PRO-10   |
| Hersteller               | GeneTex   |
| Kategorie                | Proteine/Peptide  |
| Applikation              | FA  |
| Spezies Reaktivität      | Human   |
| Konjugation              | Unconjugated  |
| NCBI                     | <a href="#">1048</a>  |
| UniProt                  | <a href="#">P06731</a>  |
| Puffer                   | Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300. |
| Expression System        | E. coli   |
| Formulierung             | Lyophilized powder  |
| Sequenz                  | N-terminal His-Tag, Ala566~Gly698 (NP_001278413.1)  |

|                        |   |
|------------------------|---|
| Anwendungsbeschreibung | <p>Carcinoembryonic antigen (CEA) which are characterized as members of the CD66 cluster of differentiation have highly related glycoproteins involved in cell adhesion. It also can be used as a tumor marker in clinical tests. CEA are glycosyl phosphatidyl inositol (GPI) cell-surface-anchored glycoproteins whose specialized sialofucosylated glycoforms serve as functional colon carcinoma L-selectin and E-selectin ligands. Besides, Galectin 4 (GAL4) has been identified as an interactor of CEA, thus a binding ELISA assay was conducted to detect the interaction of recombinant human CEA and recombinant human GLA4. Briefly, CEA were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to GLA4-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-CEA pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of CEA and GLA4 was in a dose dependent manner.</p> |
|------------------------|---|