

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Human IL1 Receptor antagonist protein, His tag, Unconjugated GTX00277-PRO

Artikelname	Human IL1 Receptor antagonist protein, His tag, Unconjugated
Artikelnummer	GTX00277-PRO
Hersteller Artikelnummer	GTX00277-pro
Alternativnummer	GTX00277-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	<a href="#">3557</a>
UniProt	<a href="#">P18510</a>
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	HEK293 cells
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Arg26~Glu177 (NP_000568.1)

Anwendungsbeschreibung

IL-1RA is an agent that binds non-productively to the cell surface interleukin-1 receptor (IL-1R), the same receptor that binds interleukin 1 (IL-1), preventing IL-1 from sending a signal to that cell. Besides, Interleukin 1 Alpha (IL-1a) has been identified as an interactor of IL-1RA, thus a binding ELISA assay was conducted to detect the interaction of recombinant human IL-1RA and recombinant human IL-1a. Briefly, IL1RA were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to IL-1a-coated microtiter wells and incubated for 2 h at 37C. Wells were washed with PBST and incubated for 1 h with anti-IL-1RA pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of IL-1RA and IL-1a was in a dose dependent manner.