

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Mouse PF4 protein, His tag (active), Unconjugated GTX00334-PRO

Artikelname	Mouse PF4 protein, His tag (active), Unconjugated
Artikelnummer	GTX00334-PRO
Hersteller Artikelnummer	GTX00334-pro
Alternativnummer	GTX00334-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Mouse
Konjugation	Unconjugated
NCBI	<a href="#">56744</a>
UniProt	<a href="#">Q9Z126</a>
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Val30~Ser105 (NP_064316.1)

Anwendungsbeschreibung

Platelet factor 4 (PF4) is a small cytokine belonging to the CXC chemokine family that is also known as chemokine (C-X-C motif) ligand 4 (CXCL4). This chemokine is released from alpha-granules of activated platelets during platelet aggregation, and promotes blood coagulation by moderating the effects of heparin-like molecules. Due to these roles, it is predicted to play a role in wound repair and inflammation. To measure its ability to inhibit the FGF basic-dependent proliferation of HUVEC human umbilical vein endothelial cells, HUVEC cells were seeded into 96-well plates at a density of 3000 cells/well with 2% serum standard DMEM including 1µg/ml recombinant human FGF1 and various concentrations of recombinant human PF4. After incubated for 48h, cells were observed by inverted microscope. Briefly, 10 µl of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-2 hours at 37C. Proliferation of HUVEC cells after incubation with PF4 for 48h observed by inverted microscope.