

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Rat Factor VII protein, His tag, Unconjugated GTX00352-PRO

Artikelname	Rat Factor VII protein, His tag, Unconjugated
Artikelnummer	GTX00352-PRO
Hersteller Artikelnummer	GTX00352-pro
Alternativnummer	GTX00352-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Rat
Konjugation	Unconjugated
NCBI	<a href="#">260320</a>
UniProt	<a href="#">Q8K3U6</a>
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Ala315~Asp433 (NP_690059.1)

Anwendungsbeschreibung

The main role of factor VII (FVII) is to initiate the process of coagulation in conjunction with tissue factor (TF). Tissue factor is found on the outside of blood vessels-normally not exposed to the bloodstream. Upon vessel injury, tissue factor is exposed to the blood and circulating FVII. Once bound to TF, FVII is activated to FVIIa by different proteases, among which are thrombin (factor IIa), factor Xa, IXa, XIIa, and the FVIIa-TF complex itself. Tissue Factor (TF) has been identified as an interactor of FVII, thus a binding ELISA assay was conducted to detect the interaction of recombinant rat FVII and recombinant rat TF. Briefly, FVII were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to TF-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-FVII pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of of FVII and TF was in a dose dependent manner.