

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-CD20 Antibody, Clone: [SPM618], Mouse, Monoclonal NSJ-V2390IHC-7ML

Artikelname	Anti-CD20 Antibody, Clone: [SPM618], Mouse, Monoclonal
Artikelnummer	NSJ-V2390IHC-7ML
Hersteller Artikelnummer	V2390IHC-7ML
Alternativnummer	NSJ-V2390IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant human MS4A1/CD20 protein was used as the immunogen for this anti-CD20 antibody.
Produktbeschreibung	This antibody recognizes a protein of 33-37kDa, identified as CD20. The epitope is similar to or identical to that recognized by other CD20 antibodies including Leu-16 and B1. This antibody can be used for immunophenotyping of leukemia and malignant ...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM618]
UniProt	P11836
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the anti-CD20 antibody to be titrated up or down for optimal performance. 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.