

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

LCA Antibody, Clone: [SPM496], Mouse, Monoclonal NSJ-V2400-100UG

Artikelname	LCA Antibody, Clone: [SPM496], Mouse, Monoclonal
Artikelnummer	NSJ-V2400-100UG
Hersteller Artikelnummer	V2400-100UG
Alternativnummer	NSJ-V2400-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Stimulated human leukocytes were used as the immunogen for the LCA antibody.
Produktbeschreibung	CD45R, also designated CD45, B220, LCA and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Along with other members of the PTP family, it regulates a number of cellular processes including cell...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM496]
UniProt	P08575
Puffer	1X PBS, pH 7.4

Reinheit	Protein G affinity chromatography
Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Due to variation in protocol and secondary antibody used, the LCA antibody may need to be titrated for optimal performance. 1. FFPE staining requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.