

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytochrome C Antibody, Clone: [CYCS/1010], Mouse, Monoclonal NSJ-V2789-100UG

Artikelname	Cytochrome C Antibody, Clone: [CYCS/1010], Mouse, Monoclonal
Artikelnummer	NSJ-V2789-100UG
Hersteller Artikelnummer	V2789-100UG
Alternativnummer	NSJ-V2789-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human protein was used as the immunogen for the Cytochrome C antibody.
Produktbeschreibung	Cytochrome C, or CYCS, is an electron carrier protein. The oxidized form of the CYCS heme group can accept an electron from the heme group of the Cytochrome C1 subunit of cytochrome reductase. CYCS then transfers this electron to the cytochrome oxida...
Klonalität	Monoclonal
Klon-Bezeichnung	[CYCS/1010]
UniProt	P99999
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Western blot: 1-2ug/ml,Immunohistochemistry (FFPE): 0.25-0.5ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the Cytochrome C antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.