

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Pan Cytokeratin Antibody, Clone: [KRTL/1077 + KRTH/1076], Mouse, Monoclonal NSJ-V3051IHC-7ML**

|                          |   |
|--------------------------|---|
| Artikelname              | Pan Cytokeratin Antibody, Clone: [KRTL/1077 + KRTH/1076], Mouse, Monoclonal   |
| Artikelnummer            | NSJ-V3051IHC-7ML  |
| Hersteller Artikelnummer | V3051IHC-7ML  |
| Alternativnummer         | NSJ-V3051IHC-7ML  |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC-P   |
| Spezies Reaktivität      | Human, Rat  |
| Immunogen                | Recombinant human KRT77 and KRT76 protein were used as the immunogen for the pan Cytokeratin antibody.  |
| Produktbeschreibung      | Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, which include Keratins 1, 3, 4, 5, 6, 8, 1... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [KRTL/1077 + KRTH/1076]   |
| UniProt                  | pan   |

|                        |  |
|------------------------|--|
| Reinheit               | Protein G affinity chromatography  |
| Formulierung           | Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*   |
| Antibody Type          | Primary Antibody   |
| Application Verdünnung | The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.  |
| Anwendungsbeschreibung | Optimal dilution of the pan Cytokeratin antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |