

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Pan Cytokeratin Antibody, Clone: [KRTL/1077 + KRTH/1076], Mouse, Monoclonal NSJ-V3051SAF-100UG

| | |
|--------------------------|---|
| Artikelname | Pan Cytokeratin Antibody, Clone: [KRTL/1077 + KRTH/1076], Mouse, Monoclonal |
| Artikelnummer | NSJ-V3051SAF-100UG |
| Hersteller Artikelnummer | V3051SAF-100UG |
| Alternativnummer | NSJ-V3051SAF-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human, Rat |
| Immunogen | Recombinant human KRT77 and KRT76 protein were used as the immunogen for the pan Cytokeratin antibody. |
| Produktbeschreibung | Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, which include Keratins 1, 3, 4, 5, 6, 8, 1... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [KRTL/1077 + KRTH/1076] |
| UniProt | pan |

| | |
|------------------------|--|
| Reinheit | Protein G affinity chromatography |
| Formulierung | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the pan Cytokeratin antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |