

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Blood Group Lewis a Antibody, Clone: [7LE], Mouse, Monoclonal NSJ-V3058IHC-7ML

Artikelname	Blood Group Lewis a Antibody, Clone: [7LE], Mouse, Monoclonal
Artikelnummer	NSJ-V3058IHC-7ML
Hersteller Artikelnummer	V3058IHC-7ML
Alternativnummer	NSJ-V3058IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human, Mouse
Immunogen	Mucins isolated from ovarian cyst fluid were used as the immunogen for the Blood Group Lewis a antibody.
Produktbeschreibung	Recognizes a carbohydrate determinant of Gal 1-3 (Fuc 1-4) GlcNAc which is blood group antigen Lewis A. It is present primarily on epithelial cells such as colon and kidneys. In the tumors and dedifferentiated tissues, decrease of Lewis A antigen was...
Klonalität	Monoclonal
Klon-Bezeichnung	[7LE]
UniProt	Not Applicable
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the Blood Group Lewis a antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.