

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **hCG Holo Antibody, Clone: [HCGab/52], Mouse, Monoclonal NSJ-V3064-100UG**

Artikelname	hCG Holo Antibody, Clone: [HCGab/52], Mouse, Monoclonal
Artikelnummer	NSJ-V3064-100UG
Hersteller Artikelnummer	V3064-100UG
Alternativnummer	NSJ-V3064-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Purified hCG protein was used as the immunogen for the hCG Holo antibody.
Produktbeschreibung	This mAb reacts only with the intact-HCG and not with either free alpha- or free beta-chain of HCG. HCG is a glycoprotein and is composed of two non-identical, non-covalently linked polypeptide chains designated as the alpha and beta subunits. The al...
Klonalität	Monoclonal
Klon-Bezeichnung	[HCGab/52]
UniProt	<a href="#">P01215</a>
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the hCG Holo antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.