

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Pan Muscle Actin Antibody Cocktail, Clone: [HHF35 + MSA/953], Mouse, Monoclonal NSJ-V3102-20UG

Artikelname	Pan Muscle Actin Antibody Cocktail, Clone: [HHF35 + MSA/953], Mouse, Monoclonal
Artikelnummer	NSJ-V3102-20UG
Hersteller Artikelnummer	V3102-20UG
Alternativnummer	NSJ-V3102-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human, Rat
Immunogen	Recombinant human actin fragment was used as the immunogen for the pan Muscle Actin antibody cocktail.
Produktbeschreibung	This antibody recognizes actin of skeletal, cardiac, and smooth muscle cells. It is not reactive with other mesenchymal cells except for myoepithelium. Actin can be resolved on the basis of its isoelectric points into three distinctive components: al...
Klonalität	Monoclonal
Klon-Bezeichnung	[HHF35 + MSA/953]
UniProt	P62736

Reinheit	Protein G affinity chromatography
Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 0.25-0.5ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the pan Muscle Actin antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.