

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Double Stranded DNA Antibody, Clone: [DSD/958], Mouse, Monoclonal NSJ-V3104-100UG

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Double Stranded DNA Antibody, Clone: [DSD/958], Mouse, Monoclonal                                                                                                                                                                                             |
| Artikelnummer            | NSJ-V3104-100UG                                                                                                                                                                                                                                               |
| Hersteller Artikelnummer | V3104-100UG                                                                                                                                                                                                                                                   |
| Alternativnummer         | NSJ-V3104-100UG                                                                                                                                                                                                                                               |
| Hersteller               | NSJ Bioreagents                                                                                                                                                                                                                                               |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC-P                                                                                                                                                                                                                                                         |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Nuclei of Burkitts cells were used as the immunogen for the Double Stranded DNA antibody.                                                                                                                                                                     |
| Produktbeschreibung      | This mAb recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. This mAb produces a homogeneous staining pattern in the nucleus of norma... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Klon-Bezeichnung         | [DSD/958]                                                                                                                                                                                                                                                     |
| UniProt                  | Not Known                                                                                                                                                                                                                                                     |
| Reinheit                 | Protein G affinity chromatography                                                                                                                                                                                                                             |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Formulierung           | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide                                                                                                                                                                                                                                                                                                                                                                                           |
| Antibody Type          | Primary Antibody                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT                                                                                                                                                                                                                                                                                                                                                                                                               |
| Anwendungsbeschreibung | Optimal dilution of the Double Stranded DNA antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |