

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Double Stranded DNA Antibody, Clone: [DSD/958], Mouse, Monoclonal NSJ-V3104-20UG

Artikelname	Double Stranded DNA Antibody, Clone: [DSD/958], Mouse, Monoclonal
Artikelnummer	NSJ-V3104-20UG
Hersteller Artikelnummer	V3104-20UG
Alternativnummer	NSJ-V3104-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Nuclei of Burkitts cells were used as the immunogen for the Double Stranded DNA antibody.
Produktbeschreibung	This mAb recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. This mAb produces a homogeneous staining pattern in the nucleus of norma...
Klonalität	Monoclonal
Klon-Bezeichnung	[DSD/958]
UniProt	Not Known
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the Double Stranded DNA antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.