

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Basic Cytokeratin Antibody, Clone: [SPM591], Mouse, Monoclonal NSJ-V3106-20UG

Artikelname	Basic Cytokeratin Antibody, Clone: [SPM591], Mouse, Monoclonal
Artikelnummer	NSJ-V3106-20UG
Hersteller Artikelnummer	V3106-20UG
Alternativnummer	NSJ-V3106-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Solubilized keratin extract from human stratum corneum was used as the immunogen for the Basic Cytokeratin antibody.
Produktbeschreibung	This antibody recognizes CK1, CK5, CK10 and CK14. In normal epithelia, it stains stratified epithelia, myoepithelial cells and basal cells in the prostate gland and bronchi. This mAb shows no reactivity with hepatocytes, pancreatic acinar cells, prox...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM591]
UniProt	Q01546
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the Basic Cytokeratin antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.