

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MUC1 Antibody Cocktail, Clone: [GP1.4 + E29], Mouse, Monoclonal NSJ-V3138IHC-7ML

Artikelname	MUC1 Antibody Cocktail, Clone: [GP1.4 + E29], Mouse, Monoclonal
Artikelnummer	NSJ-V3138IHC-7ML
Hersteller Artikelnummer	V3138IHC-7ML
Alternativnummer	NSJ-V3138IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Human milk fat globule membranes were used as the immunogen for this MUC1 antibody.
Produktbeschreibung	In Western blotting, it recognizes proteins in MW range of 265-400kDa, identified as different glycoforms of EMA. The alpha subunit has cell adhesive properties. It can act both as an adhesion and an anti-adhesion protein. EMA may provide a protectiv...
Klonalität	Monoclonal
Klon-Bezeichnung	[GP1.4 + E29]
UniProt	P15941
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	The optimal dilution of the MUC1 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.