

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Chromogranin A Antibody Cocktail, Clone: [LK2H10 + PHE5], Mouse, Monoclonal NSJ-V3140-20UG

Artikelname	Chromogranin A Antibody Cocktail, Clone: [LK2H10 + PHE5], Mouse, Monoclonal
Artikelnummer	NSJ-V3140-20UG
Hersteller Artikelnummer	V3140-20UG
Alternativnummer	NSJ-V3140-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Human pheochromocytoma cells were used as the immunogen for this Chromogranin A antibody cocktail (LK2H10 & PHE5).
Produktbeschreibung	Chromogranin A is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, pheochromocytomas, para...
Klonalität	Monoclonal
Klon-Bezeichnung	[LK2H10 + PHE5]
UniProt	P10645

Reinheit	Protein G affinity chromatography
Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 0.1-0.2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the Chromogranin A antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.