

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

DOG-1 Antibody Cocktail, Clone: [DG1/447 + DOG-1.1], Mouse, Monoclonal NSJ-V3159IHC-7ML

Artikelname	DOG-1 Antibody Cocktail, Clone: [DG1/447 + DOG-1.1], Mouse, Monoclonal
Artikelnummer	NSJ-V3159IHC-7ML
Hersteller Artikelnummer	V3159IHC-7ML
Alternativnummer	NSJ-V3159IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant human DOG1 protein (DG1/447) and amino acids 904-986 (DOG-1.1) were used as the immunogens for this DOG-1 antibody cocktail.
Produktbeschreibung	Expression of DOG-1 protein is elevated in the gastrointestinal stromal tumors (GISTs), c-kit signaling-driven mesenchymal tumors of the GI tract. DOG-1 is rarely expressed in other soft tissue tumors, which, due to appearance, may be difficult to di...
Klonalität	Monoclonal
Klon-Bezeichnung	[DG1/447 + DOG-1.1]
UniProt	Q5XXA6

Reinheit	Protein G affinity chromatography
Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	The optimal dilution of the DOG-1 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.