

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

DOG-1 Antibody Cocktail, Clone: [DG1/447 + DOG-1.1], Mouse, Monoclonal NSJ-V3159SAF-100UG

Artikelname	DOG-1 Antibody Cocktail, Clone: [DG1/447 + DOG-1.1], Mouse, Monoclonal
Artikelnummer	NSJ-V3159SAF-100UG
Hersteller Artikelnummer	V3159SAF-100UG
Alternativnummer	NSJ-V3159SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant human DOG1 protein (DG1/447) and amino acids 904-986 (DOG-1.1) were used as the immunogens for this DOG-1 antibody cocktail.
Produktbeschreibung	Expression of DOG-1 protein is elevated in the gastrointestinal stromal tumors (GISTs), c-kit signaling-driven mesenchymal tumors of the GI tract. DOG-1 is rarely expressed in other soft tissue tumors, which, due to appearance, may be difficult to di...
Klonalität	Monoclonal
Klon-Bezeichnung	[DG1/447 + DOG-1.1]
UniProt	Q5XXA6

Reinheit	Protein G affinity chromatography
Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the DOG-1 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.