

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Mucin-1 Antibody / Rabbit Monoclonal, Clone: [MUC1/1887R] NSJ-V3740-20UG

| | |
|--------------------------|---|
| Artikelname | Recombinant Mucin-1 Antibody / Rabbit Monoclonal, Clone: [MUC1/1887R] |
| Artikelnummer | NSJ-V3740-20UG |
| Hersteller Artikelnummer | V3740-20UG |
| Alternativnummer | NSJ-V3740-20UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC-P, WB |
| Spezies Reaktivität | Human |
| Immunogen | Human milk fat globule membranes were used as the immunogen for this recombinant Mucin-1 antibody. |
| Produktbeschreibung | Mucin-1/MUC1 protein is encoded by the MUC1 gene in humans. This gene encodes a membrane-bound protein that is a member of the mucin family. Family members are O-glycosylated proteins that play an essential role in forming protective mucous barriers ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [MUC1/1887R] |
| UniProt | P15941 |
| Reinheit | Protein A affinity chromatography |

| | |
|------------------------|---|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT,Western blot: 1-2ug/ml |
| Anwendungsbeschreibung | The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the recombinant Mucin-1 antibody to be titered up or down for optimal performance.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |